



PROOF OF PURCHASE
COMPROBANTE DE COMPRA

Preen®

LAWN

STEP SAVER™ Weed Control PLUS FERTILIZER

Control de malezas y fertilización
de césped en menos pasos

Stops More Weeds In One Application!

- One product kills broadleaf weeds and prevents crabgrass
- Fertilizes longer than most other leading brands
- Apply once in spring when weeds are growing



Dandelions

Crabgrass

COVERS
5,000 SQ. FT.
Net Wt. 18 lbs. (8.16 kg)

KEEP OUT OF REACH OF CHILDREN
CAUTION

See back panel for Precautionary Statements.

26-2-8

ACTIVE INGREDIENTS:	
2,4-D.....	0.64%
Mecoprop-p.....	0.14%
Dicamba.....	0.06%
Dithiopyr.....	0.16%
OTHER INGREDIENTS:.....	99.00%
TOTAL.....	100.00%

THIS PRODUCT CONTAINS:
12.80 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton.
2.80 lbs. (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per ton.
1.20 lbs. 3,6-dichloro-o-anisic acid equivalent per ton.
3.20 lbs. dithiopyr per ton.

Isomer Specific by AOAC Methods
Contains the single isomer form of Mecoprop-p.

Stops More Weeds In One Application!

LAWN

STEP SAVER™ Weed Control PLUS FERTILIZER

Control de malezas y fertilización
de césped en menos pasos

Stops More Weeds In One Application!

- One product kills broadleaf weeds and prevents crabgrass
- Fertilizes longer than most other leading brands
- Apply once in spring when weeds are growing



Dandelions

Crabgrass

COVERS
5,000 SQ. FT.
Net Wt. 18 lbs. (8.16 kg)

KEEP OUT OF REACH OF CHILDREN
CAUTION
See back panel for Precautionary Statements.

26-2-8

ACTIVE INGREDIENTS:	
2,4-D	0.64%
Mecoprop-p	0.14%
Dicamba	0.06%
Dithiopyr	0.16%
OTHER INGREDIENTS:	99.00%
TOTAL	100.00%

THIS PRODUCT CONTAINS:
12.80 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton.
2.80 lbs. (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per ton.
1.20 lbs. 3,6-dichloro-o-anisic acid equivalent per ton.
3.20 lbs. dithiopyr per ton.

Isomer Specific by AOAC Methods
Contains the single isomer form of Mecoprop-p.

NOTICE: Read the entire label. Use only according to label directions.
STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL PRECAUTIONS AND RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.
- If incorporation is accomplished by watering-in, do not enter or allow entry to the treated areas (except those involved in watering) until the watering-in is complete and the surface is dry.
- Read all precautions and directions for use carefully before applying this product.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Keep in original container in a cool, dry area, away from extreme heat and direct sunlight. Do not store near food feed, seeds or other pesticides.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

General Information

Preen Lawn STEPSAVER Weed Control Plus Fertilizer will control all your lawn's weeds in one application. It will kill over 200 lawn weeds commonly found in home lawns including dandelion, chickweed, thistle and clover, and will prevent crabgrass and other undesirable annual grasses. Preen Lawn Step Saver Weed Control Plus Fertilizer also contains a long-lasting, timed release fertilizer that has all the nutrients your lawn needs for 2 to 3 months, depending on grass variety.

Preen Lawn STEPSAVER Weed Control Plus Fertilizer is recommended for the following established turf species:

Established Cool Season Turfgrasses — Kentucky bluegrass, perennial ryegrass, fine and tall fescues.

Established Warm Season Turfgrasses — Bahiagrass, bermudagrass, buffalograss, centipede grass, zoysiagrass, St. Augustinegrass (except 'Floritam').

Do not use this product on seedling buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots.

Do not use Preen Lawn STEPSAVER Weed Control Plus Fertilizer on colonial bentgrass, dichondra, carpetgrass, or 'Floritam' St. Augustinegrass.

If the lawn has been seeded recently, wait until after the third mowing before using this product. Do not seed, overseed or sprig the treated area for 10 weeks after application.

GENERAL USE PRECAUTIONS AND RESTRICTIONS

Use of Preen Lawn STEPSAVER Weed Control Plus Fertilizer on lawns and ornamental turfgrasses that are not well established, or have been weakened by stress due to mechanical operations, pests, excessive temperatures, or lack of moisture may increase the chances for turfgrass injury. Allow turfgrass weakened by these stresses to recover before application of this product.

Do not use this product where desirable clovers are present or spread (apply) the granules onto flowers, vegetables or ornamental shrubbery. Do not spread over exposed roots of trees and ornamentals. Do not exceed specified dosages for any area; be particularly careful within dripline of trees and other ornamental species.

Do not use clippings from treated areas for mulching, vegetables, or fruit trees.

Do not contaminate domestic or irrigation waters.

Do not use this product on or near desirable plants, including within the drip line of the roots of desirable trees and shrubs, since injury may result.

Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches.

Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans.


Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.



When to Apply

Preen Lawn STEPSAVER Weed Control Plus Fertilizer can be used either in the spring or fall. It should be used when the grass and weeds are actively growing and the soil is moist. Do not apply under drought conditions.

In the spring, the ideal time to apply Preen Lawn STEPSAVER Weed Control Plus Fertilizer is after the forsythia has bloomed or when the first dandelion flowers appear. In southern areas (see map), make a second application in 6-8 weeks for best weed and crabgrass control and for full season fertilization.

In the fall, apply Preen Lawn STEPSAVER Weed Control Plus Fertilizer when the weather cools and both the grass and weeds are actively growing, generally around Labor Day.



 Cool Season and Transition Zone Turfgrasses (Northern Area)
 Warm Season Turfgrasses (Southern Area)

How to Apply

This product needs to stick to the leaves of broadleaf weeds and should be applied when the grass is wet from dew or rain; poor results will occur if it is applied to dry lawns. The lawn should not be watered or disturbed for 24 hours after application. Delay mowing for 3 days after application.

Apply this product when the wind is calm. Do not apply this product onto desirable plants or shrubs as damage may result. If the product gets onto other plants immediately brush or wash it off the foliage.

Use a drop or rotary spreader designed to apply granular fertilizers. Avoid skips or excessive overlaps as poor weed control or turfgrass injury may result.

How Much to Use

Apply 3.6 lbs. of product per 1,000 sq. ft. of turfgrass.

In New York State: Do not exceed one application at the recommend rate of 3.6 lb of product per 1,000 sq. ft. (0.25 active ingredient of dithiopyr per acre).

Other States: The maximum broadcast application rate is 3.6 pounds of product per 1,000 sq.ft. per application (1.0 lb 2,4-D ae, 0.22 lb MCPP-p ae, 0.09 lb dicamba ae and 0.25 lb dithiopyr per acre per application). The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 7.2 pounds of product per 1,000 sq.ft. (2.0 lb 2,4-D ae, 0.44 lb MCPP-p ae and 0.18 lb dicamba ae and 0.50 lb dithiopyr per acre per year).

Calibration Instructions

For best results, apply this product evenly and uniformly avoiding streaking, skips or overlaps.


For 3.6 lbs. of product per 1,000 sq. ft. of turfgrass: check your spreader on 275 sq. ft. and adjust to apply at the rate of 1 lb. of product per 275 sq. ft.

Check equipment frequently to verify calibration.

Suggested Spreader Settings

These spreader settings are approximate. The age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

Broadcast Spreaders	Setting
Agri-Fab® Rotary	4
Cyclone® Rotary	3 1/2
Earthway® Rotary	14
Greenview® Jet-Spred® SS1	14



Do not contaminate domestic or irrigation waters.

Do not use this product on or near desirable plants, including within the drip line of the roots of desirable trees and shrubs, since injury may result.

Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches.

Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans.

Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

Weeds Controlled

When applied as directed, Preen Lawn STEPSAVER Weed Control Plus Fertilizer will control the following weeds:

Annual Grass weeds:

Annual Bluegrass (<i>Poa annua</i>)	Barnyardgrass	Foxtail (green and yellow)	Perennial Ryegrass
Annual Ryegrass	Crabgrass	Goosegrass	Smutgrass (seedling)
	Crowfootgrass	Kikyugrass (seedling)	

Established Broadleaf weeds:


Annual yellow sweetclover	Catnip	Hoary plantain	Prostrate knotweed	Tumble mustard
Aster	Catsear	Hoary vervain	Prostrate pigweed	Tumble pigweed
Austrian fieldcress	Chickweed:	Horsenettle	Prostrate spurge	Velvetleaf
Bedstraw	common, mouseear	Jimsonweed	Prostrate vervain	Venice mallow
Beggarticks	Chicory	Knawel	Puncturevine	Virginia buttonweed
Betony, Florida	Cinquefoil	Knotweed	Purslane, common	Virginia creeper
Bindweed, field	Clover: crimson, hop	Kochia	Ragweed	Virginia pepperweed
Bird vetch	red, strawberry,	Lambsquarters	Red sorrel	Wavyleaf bullthistle
Bitter wintercress	sweet, white	Lespedeza	Redroot pigweed	Western clematis
Bittercress, hairy	Cockle	Mallow	Redstem filaree	Western salsify
Bitterweed	Cocklebur	Matchweed	Rough cinquefoil	White mustard
Black-eyed Susan	Common mullein	Mexicanweed	Rough fleabane	Wild aster
Black medic	Creeping jenny	Milk vetch	Russian pigweed	Wild buckwheat
Black mustard	Cudweed	Morningglory	Scarlet pimpernel	Wild carrot
Blackseed plantain	Curly dock	Mouseear hawkweed	Scotch thistle	Wild four-o'clock
Blessed thistle	Daisy: English,	Muswort	Sheep sorrel	Wild garlic
Bloodflower	oxeye	Musk thistle	Shepherdspurse	Wild geranium
milkweed	Daisy fleabane	Mustard	Slender plantain	Wild lettuce
Blue lettuce	Dandelion	Narrowleaf plantain	Smallflower	Wild marigold
Blue vervain	Dichondra	Narrowleaf vetch	galinsoga	Wild mustard
Bracted plantain	Dogfennel	Nettle	Smooth dock	Wild onion
Brassbuttons	Dollarweed	Orange hawkweed	Smooth pigweed	Wild parsnip
Bristly oxtongue	False dandelion	Oriental cocklebur	Sorrel	Wild radish
Broadleaf dock	Falseflax	Oxalis	Sowthistle	Wild rape
Broadleaf plantain	False sunflower	Parsley-piert	Spanishneedles	Wild sweet potato
Broomweed	Fiddleneck	Parsnip	Speedwell	Wild vetch
Buckhorn plantain	Florida pusley	Pearlwort	Spiny amaranth	Woodsorrel
Bulbous buttercup	Frenchweed	Pennycress	Spiny cocklebur	Woolly croton
Bull thistle	Galinsoga	Pepperweed	Spiny sowthistle	Woolly morningglory
Bullnettle	Goathead	Pigweed	Spotted catsear	Woolly plantain
Burclover	Goldenrod	Pineappleweed	Spotted spurge	Wormseed
Burdock	Ground ivy	Pineywoods	Spurweed	Yarrow
Burning nettle	Gumweed	bedstraw	Stinging nettle	Yellow rocket
Burweed	Hairy fleabane	Plains tickseed	Strawberry, India	Yellowflower
Buttercup	Hawkweed	Plantain	mock	pepperweed &
Buttonweed	Healall	Poison ivy	Tall nettle	other broadleaf
Canada thistle	Heartleaf drymary	Poison oak	Tall vervain	weeds
Carolina geranium	Heath aster	Pokeweed	Tansy ragwort	
Carpetweed	Hedge bindweed	Poorjoe	Tansy mustard	
Catchweed	Hedge mustard	Prairie sunflower	Tanweed	
bedstraw	Henbit	Prickly lettuce	Thistle	
	Hoary cress	Prickly sida	Trailing crownvetch	

Guaranteed Fertilizer Analysis:


Total Nitrogen (N)	26%
3.0% Ammoniacal Nitrogen	
7.0% Water Insoluble Nitrogen*	
9.6% Urea Nitrogen	
6.4% Other Water Soluble Nitrogen*	
Available Phosphate (P ₂ O ₅)	2%
Soluble Potash (K ₂ O)	8%
Sulfur (S)	3.0%
3.0% Combined Sulfur (S)	
Derived from: Ammonium Phosphate, Ammonium Sulfate, Methylene Ureas, Urea, Muriate of Potash.	
Chlorine (Cl) not more than	7.0%
* 13.4% Slowly Available Nitrogen from Methylene Ureas.	F699

Suggested Spreader Settings

These spreader settings are approximate. The age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.



Apply in a single pass at a rate of 3.6 lbs. per 1,000 sq. ft. Start on the outside border and apply in a circular path working toward the center. Overlap the coverage slightly, and shut off spreader before making hard turns.



Apply in a single pass at a rate of 3.6 lbs. per 1,000 sq. ft. Start on the outside border and apply in a rectangular path working back and forth across the length of the lawn. Overlap the coverage slightly, and shut off spreader before making hard turns.

Broadcast Spreaders

Setting

Agri-Fab® Rotary	4
Cyclone® Rotary	3 1/2
Earthway® Rotary	14
Greenview® Jet-Spred® SS1	14
Lesco® Rotary (dial A-S)/(number dial)	H / 14
Red Devil® Rotary (1-20)	5
Republic® Rotary	4
Scotts® EasyGreen®	23
Scotts® EdgeGuard®	4 1/2
Scotts® SpeedyGreen® 2000, 3000	4 1/2
Sears® Craftsman Rotary	4
Sta-Green® Rotary (1-10)	2 1/2
True Temper® Rotary	3 1/2
Vigoro® Rotary (1-10)	2 1/2



Drop Spreaders

Setting

Agri-Fab® Drop	4
Greenview® Drop-Spred® XP1	16
Red Devil® Drop (1-5)	3 1/2
Republic® Drop	5
Scotts® Drop AccuGreen® (2-18)	6
Scotts® Drop PF1 (2-18)	6
Sta-Green® Drop (1-5)	3 1/2



If your spreader model does not appear above, go to our website at www.preen.com, or call toll-free (800) 233-1067, 8 A.M. - 5 P.M. EDT, Monday - Friday, for additional model settings.

Warranty

The manufacturer warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions not reasonably foreseeable to manufacturer, and buyer assumes the risk of any such use.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on this label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of seller. All such risks shall be assumed by buyer.

Limitation of Remedies

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

Seller shall not be liable for losses or damages resulting from handling or use of this product unless seller is promptly notified of such loss or damage in writing. In no case shall seller be liable for consequential or incidental damages or losses.

PREEN GUARANTEE

Lebanon Seaboard Corporation guarantees you will be satisfied with the performance of this product when used as directed, or your money back. If not satisfied, describe the nature of your problem, and send with proof of purchase to:

LEBANON SEABOARD CORPORATION
1600 EAST CUMBERLAND STREET, LEBANON, PA 17042



1-800-233-1067

www.preen.com



PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Harmful if absorbed through skin or if swallowed. Avoid contact with skin, eyes or clothing. Handlers should wear long sleeved shirt and long pants, shoes plus socks, and chemical-resistant gloves made of any waterproof material.

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent material that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them.

User Safety Recommendations

- Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

EPA Reg. No. 2217-859-961

(L) (D)

EPA Est. 961-PA-01:961-IL-01

Superscript used is first letter of lot number.

CAS Registry Numbers:

2,4-D CAS No. 94-75-7

Mecoprop-p CAS No. 16484-77-8

Dicamba CAS No. 1918-00-9

Dithiopyr CAS No. 97886-45-8

Preen® is a registered trademark
of Lebanon Seaboard Corporation.

Information regarding the contents and levels of
metals in this product is available on the Internet
at <http://www.regulatory-info-lebsea.com> 7/05



PROOF OF PURCHASE
COMPROBANTE DE COMPRA

FIRST AID

IF IN EYES:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.• Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For additional information in case of emergency, call toll free (888) 208-1368.	